

Morphology of three marine epizoic diatoms from the Mexican Pacific: *Protoraphis hustediana*, *Pseudohimantidium pacificum* and *Sceptronema orientale* (Bacillariophyta)

Morfología de tres diatomeas epizoicas marinas del Pacífico mexicano: *Protoraphis hustediana*, *Pseudohimantidium pacificum* y *Sceptronema orientale* (Bacillariophyta)

David U. Hernández-Becerril

Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, México

*Autor de correspondencia: dhernand@cmarl.unam.mx

Recibido: 11 de junio de 2021

Aceptado: 27 de septiembre 2021

Publicación en línea: xxxx

Abstract

Key words:
diatoms; electron microscopy;
morphological characters; new records,
tychoplankton

Diatoms are a successful taxonomic group of microbes, which exhibit a high diversity in marine habitats. Marine epibiont diatoms also display an important diversity of symbiotic associations. Net phytoplankton samples collected in the Mexican Pacific were analyzed, and apart from finding planktonic species, three epizoic diatom species were additionally observed: *Protoraphis hustediana*, *Pseudohimantidium pacificum* and *Sceptronema orientale*. The three species, usually found attached to copepods, were studied by light, scanning and transmission electron microscopy (ML, SEM and TEM), showing morphological characters such as shape of the valves, central (and sigmoid) sternum, fine striae, areolae shape, apical and aggregated rimoportulae, and slit fields at the apices (or poles). Morphological details found in *Pseudohimantidium pacificum* and in *Sceptronema orientale* include the type of velum in the areolae (rota type) and the external openings of the rimoportulae, respectively. *Protoraphis hustediana* and *Sceptronema orientale* are new records for the Mexican Pacific.

Palabras clave:
caracteres morfológicos;
diatomeas; microscopía electrónica;
nuevos registros; ticoplancton

Resumen

Las diatomeas constituyen un grupo taxonómico de microbios exitosos, que exhiben una alta diversidad en hábitats marinos. Las diatomeas epibiontes marinas también despliegan una diversidad importante de asociaciones simbióticas. Se analizaron muestras de red de fitoplancton de localidades del Pacífico mexicano, que estuvieron representadas por especies planctónicas, y adicionalmente se observaron tres especies de diatomeas epizoicas: *Protoraphis hustediana*, *Pseudohimantidium pacificum* y *Sceptronema orientale*. Las tres especies, usualmente adheridas a copépodos, fueron estudiadas mediante microscopía de luz, electrónica de barrido y transmisión (ML, MEB y MET), cuyas observaciones mostraron caracteres morfológicos como forma de las valvas, sternum central (y sigmoideo), delicada estriación, forma de las areolas, rimopórtulas agregadas en los ápices, y campos apicales (o en los polos) de ranuras, con lo que se confirman estudios previos. Algunos detalles morfológicos encontrados en *Pseudohimantidium pacificum* y *Sceptronema orientale* incluyen el tipo de velum de las areolas (del tipo rota) y las aperturas externas de las rimopórtulas, respectivamente. *Protoraphis hustediana* y *Sceptronema orientale* son nuevos registros para el Pacífico mexicano.

Diatoms belong to a very successful taxonomic group of microbes in aquatic environs all over the world (Round *et al.*, 1990; Medlin and Kaczmarek, 2004). Marine diatoms exhibit a remarkable diversity in terms of sizes, shapes and life histories, and planktonic diatoms are a very important contributor to the biomass and productivity of world's oceans (Kooistra *et al.*, 2007). Littoral, benthic and epibiont marine diatoms have been also studied, although they display a high morphologic

complexity (Round, 1971; Round *et al.*, 1990). Various diatom species use to live as epizoic forms on small planktonic animals as copepods and others, showing particular ways of attachment (*e.g.* developing stalks or peduncles, producing mucilage, etc.) (Hiromi and Takano, 1983; Tiffany, 2011; Totti *et al.*, 2011; Gárate-Lizárraga and Esqueda-Escarcega, 2018). During several oceanographic expeditions carried out along coasts of the Mexican Pacific, the high diversity of planktonic

diatoms has been confirmed (Hernández-Becerril *et al.*, 2010, 2013), but additionally epipsammic forms have been also reported, including descriptions of two new species (Hernández-Becerril and Barón-Campis, 2008; Hernández-Becerril *et al.*, 2012).

Three species of araphid diatoms, *Protoraphis hustediana* Simonsen, *Pseudohimantidium pacificum* Hustedt et Krasske and *Sceptronema orientale* Takano, are considered as epizoic, living on certain species of copepods. There are some modern studies of these three species, especially showing the way they attach to the copepods, some morphologic details (including the formation of chains), and their biological affinities (symbiosis), distributions and relative abundances (Simonsen, 1970; Gibson, 1978; Hiromi and Takano, 1983; Rivera *et al.*, 1986; Skovgaard and Saiz, 2006; Gómez *et al.*, 2018, 2020; Li *et al.*, 2020).

The main purpose of this work was to study the detailed morphology of these three species to contribute to the knowledge of the diversity of marine diatoms in the Mexican Pacific littorals.

This study is based on material collected during various oceanographic surveys along coasts of the Mexican Pacific during the period 1985-2019. Main spots were located at: (1) 18° 12.943' N, 103° 20.094' W, (2) 17° 47.729' N, 102° 09.294' W, (3) 17° 08.293' N, 100° 54.078' W, (4) 16° 39.310' N, 99° 55.901' W. Net samples (mesh 64 and 54 µm) were obtained by vertical hauls, the hauling depths depending on the depth of the fixed stations; a set of samples were fixed with formalin (4 %) and another set with ethanol (70 %).

Either raw or rinsed (with distilled water) samples were analyzed by Light Microscopy (LM, Olympus BX 40, attached camera Hitachi KP-D50 Color digital), where preliminary identifications, measurements and microphotographs were made. Whole rinsed samples or isolated specimens were additionally studied by Scanning Electron Microscopy (SEM, JEOL JSM6360LV) after air-drying and coating with gold. Only cleaned material was observed by Transmission Electron Microscopy (TEM, JEOL TEM1200 EXII), once the acid treatment, recommended by Hasle (1978), was followed.

Terminology for diatoms and the genera *Pseudohimantidium*, *Protoraphis* and *Sceptronema* follows that recommended by Round *et al.* (1990) and Hasle and Syvertsen (1997).

Preliminary microscopical analysis yielded a number of planktonic forms of diatoms, and also the species studied here, considered to be epizoic species, although they were not found

attached to copepods or other small animals.

Description of species

***Protoraphis hustediana* Simonsen (figures 1-4)**

References

Simonsen, 1970, p. 383, pl. 1; Hallegraeff and McWilliam, 1990, p. 39, figures 1-13; Li *et al.*, 2020, p. 121, figures 1 a-f, 2 a-f, 3 a-e.

The cells are commonly found adhered to copepods (as in the precedent species). Valves are lanceolate (slightly curved), diagonally symmetrical, with rounded apices (figure 1). The valve is convex, curving into the mantle (figure 1). The sternum is conspicuous, linear to slightly sigmoid, especially at the apices, where it becomes slightly bent at opposite sides of the rimoportulae (figure 1, 2). The striae are uniseriate, perpendicular to the main axis, and formed by rounded areolae (figure 2). There is a deep perpendicular groove near each apex of the valve, at the end of the sternum, with four external openings, representing the rimoportulae (figure 1, 3, 4). Striae are radial at the end of the valve, and apical slits fields are also apparent (figures 3, 4).


Measurements: Apical axis 45-60 µm, transapical axis 6-10 µm, 30-32 striae in 10 µm, 3-4 areolae in 1 µm.

***Pseudohimantidium pacificum* Hustedt et Krasske (figures 5-8)**


References

Krasske, 1941, p. 272, pl. 5, figure. 8; Simonsen, 1970, p. 385, figures 5-22; Gibson, 1979a, p. 149, figure 1-13; Rivera *et al.*, 1986, p. 21, figure 1-35; Round *et al.*, 1990, p. 446, figure a-g; Fernandes and Calixto-Feres, 2012, p. 837, figures 1-17; Sunesen *et al.*, 2015, p. 268, figure 23-40.


Living cells have been usually found attached to copepods by mucilage stalks produced at the apices of the frustules. In this study only loose valves were encountered, which could be positively identified as all morphological characters fit well with previous descriptions. Valves are arcuate with subrostrate apices (figure 5, 6). The valve is slightly convex and curves into the mantle. There is a curved, slightly sigmoid, and narrow sternum, at one apex curving more abruptly, ending in the apical rimoportulae and at the other apex ending in the other side of the rimoportulae (figure 5, 6). The striae are uniseriate, perpendicular to the main axis, and becoming radiate toward the apices (figure 6); areolae have a rota type (figure 8). The rimoportulae are located at the apices of the valves, with 3 to 6 rimoportulae aligned next to the other, orientated to the ventral side of the valve (figure 5, 7).



Figures 1-4. *Protoraphis hustediana*. Fig. 5. Lanceolate valve with rimoportulae (arrows), SEM. Figure 6. Detail of the central part of the valve, showing sternum and striae, SEM. Fig. 7. Detail of the apical groove, with the external opening of the rimoportulae, SEM. Figure 8. Another view of the apical groove, SEM.



Figures 5-8. *Pseudohimantidium pacificum*. Fig. 5. A complete valve showing the central sternum and the rimoportulae (arrows), LM. Figure 6. Another valve with the sternum, striae and rimoportulae (arrow), TEM. Figure 7. Detail of the rimoportulae, TEM. Figure 8. Detail of the areolae, TEM.



Figures 9-14. *Sceptronema orientale*. Figure. 9. Solitary cell with a distinct valve shape, LM. Figure. 10. Complete frustule, exhibiting the central sternum (arrow), SEM. Figure. 11. Another frustule, slightly broken at the head-pole, SEM. Figures 12, 13. Details of the head-pole, with the slit field, SEM. Figure. 14. Foot-pole showing the sternum, areolae and the external opening of a rimoportula (arrow), SEM.

***Sceptronema orientale* Takano (Figures 9-14)**

References

Takano, 1983, p. 26, figures 2, 15-20; Sar and Sunesen, 2014, p. 3, figures 2-28; Gómez *et al.*, 2020, p. 26, figures 2 A-U. Complete, living cells have been detected joined by a

mucilaginous pad to copepods. Solitary cells were found in this study (Figures 9-11). Cells in valve view have a distinct shape of an Egyptian sarcophagus (Figures 9-11). Valves are distinctly heteropolar, narrowly obovate, with a head-pole and a narrow capitate foot-pole (Figures 9-11). The valves are flat, and the sternum is central and straight (Fig. 10). The striae are uniseriate,

and are perpendicular to the sternum (Figures 10-12), whereas the areolae are round to elliptic (Figures 14). Apical slits with longitudinal bars are present at both poles, although this slit field is larger at the head-pole (Figures 12-14). These fields are most possible from where the mucilage pads emerge to adhere to copepods. There are numerous rimoportulae close to the poles, especially the foot-pole, where their elliptical external openings are visible (Figures 14).

Measurements: Apical axis 31-48 μm , transapical axis 8-10 μm , 38-42 striae in 10 μm , 4-5 areolae in 1 μm .

In this study some morphological details of three epizoic marine diatoms found in plankton samples from the Mexican Pacific are provided. These species usually live attached to copepods, although they were encountered as solitary cells or valves, most probably caught accidentally in phytoplankton net samples.

The three species studied here, *Protoraphis hustediana*, *Pseudohimantidium pacificum*, and *Sceptronema orientale* are all araphid diatoms, which usually produce mucilage pads or stalks to adhere to copepods. There are reports of large infestations of these species (Rivera *et al.*, 1986; Fernandes and Calixto-Feres, 2012; Sar and Sunesen, 2014; Gómez *et al.*, 2020). They are well adapted to the epizoic life, developing shapes and mechanisms to deal with the supposed turbulent habitat. They do share some morphological characteristics such as the presence of a central sternum (instead of a raphe) and specialized rimoportulae in aggregations or large numbers (Simonsen, 1970; Gibson, 1978; Hiromi and Takano, 1983; Rivera *et al.*, 1986; Gómez *et al.*, 2020; Li *et al.*, 2020).

The morphological characters described here have been previously shown in the literature, particularly the studies using electron microscopy, and this is a confirmation of some details, such as the shape of the valves, the central (and sigmoid in two species) sternum, the fine striae and the shape of the areolae, the apical and aggregated rimoportulae (with certain orientation in *Pseudohimantidium pacificum*), and the slit fields at the poles (especially the species *Sceptronema orientale*) (e.g. Simonsen, 1970; Gibson, 1978; Rivera *et al.*, 1986; Fernandes and Calixto-Feres, 2012; Sar and Sunesen, 2014; Gómez *et al.*, 2020; Li *et al.*, 2020). A special morphological detail found in *Pseudohimantidium pacificum* is the type of areola, which is a velum of the rota type, not previously described.

From these species, *Pseudohimantidium pacificum* has been historically recognized and is considered widely distributed and probably the best studied of the marine epizoic diatoms, whereas *Protoraphis hustediana* and *Sceptronema orientale*

have been less reported and, consequently, less studied.

The original description of *Sceptronema orientale* included a protocol mentioning the absence of rimoportulae in this species (Takano, 1983, p. 26), although his illustrations (figures 2 C and 17) suggest the presence of various rimoportulae. Later, Sar and Sunesen (2014) provided details of these rimoportulae in the species. The external openings of the rimoportulae were also found in this study.

The species *Pseudohimantidium pacificum* has been already reported from the Mexican Pacific (Gárate-Lizárraga and Muñetón-Gómez, 2009; Gárate-Lizárraga and Esqueda-Escarcega, 2018), but the other species represent new records in the Mexican Pacific, *Protoraphis hustediana* and *Sceptronema orientale*. The recognition of these two records is not a surprise, considering the poor studies dedicated to epibiont diatoms in Mexico and the relative few records all over the world (cited for each species), and obviously widens the distribution of the species in subtropical areas of the Mexican Pacific.

Acknowledgements

Partial support for this study was provided by PAPIIT, DGAPA, UNAM (project IN296516). Coordinación de la Investigación Científica (CIC, UNAM) approved and supported the use of the R/V "El Puma" to carry out the oceanographic cruises of the series "MareaR", where samplings for part of this study were performed.

References

- Fernandes, L.F. and Calixto-Feres, M. 2012. Morphology and distribution of two epizoic diatoms (Bacillariophyta) in Brazil. *Acta Botanica Brasílica* 26: 836-841. Doi: <https://doi.org/10.1590/S0102-33062012000400012>.
- Gárate-Lizárraga, I. and Esqueda-Escarcega, G.M. 2018. *Ditrichocorycaeus anglicus* (Copepoda; Poecilostomatoida), new basibiont of *Pseudohimantidium pacificum* (Bacillariophyceae) in Bahía de La Paz, Gulf of California. *CICIMAR Oceanides* 33: 63-67. Doi: <https://doi.org/10.37543/oceanides.v33i1.223>.
- Gárate-Lizárraga, I. and Muñetón-Gómez, M.S. 2009. Primer registro de la diatomea epibionte *Pseudohimantidium pacificum* y de otras asociaciones simbióticas en el Golfo de California. *Acta Botánica Mexicana* 88: 31-45. Doi: <https://doi.org/10.21829/abm88.2009.311>.

- Gibson, R.A. 1978. *Pseudohimantidium pacificum*, an epizoic diatom new to Florida Current (Western North Atlantic Ocean). *Journal of Phycology* 14: 371-373. Doi: <https://doi.org/10.1111/j.1529-8817.1978.tb00316.x>
- Gibson, R.A. 1979a. An ultrastructure study of *Pseudohimantidium pacificum* Hust. & Krasske (Bacillariophyceae: Photoraphidaceae) with special reference to the labiate processes. *Nova Hedwigia Beihefte* 64: 147-161.
- Gibson, R.A. 1979b. *Protoraphis atlantica* sp. nov., a new marine epizoic diatom. *Bacillaria* 2: 109-126.
- Gómez F., Courcot, L. and Artigas, L.F. 2020. Observations of the diatoms *Sceptronema orientale* Takano and *Tabularia parva* (Kützting) D.M. Williams and Round on the exoskeleton of copepods in the English Channel and coastal Celtic Seas. *Cryptogamie, Algologie* 41: 25-30. Doi: <https://doi.org/10.5252/cryptogamie-algologie2020v41a4>.
- Gómez, F., Wang, L. and Lin, S. 2018. Morphology and molecular phylogeny of epizoic araphid diatoms on marine zooplankton, including *Pseudofalcula hyalina* gen. & comb. nov. (Fragilariophyceae, Bacillariophyta). *Journal of Phycology* 54: 557-570. Doi: <https://doi.org/10.1111/jpy.12760>.
- Hallegraeff, G.M. and McWilliam, P.S. 1990. The complex labiate process of the epizoic diatom *Protoraphis hustediana* Simonsen. *Nova Hedwigia, Beihefte* 100: 39-45.
- Hasle, G.R. 1978. Diatoms. In: Sournia, A., Editor. *Phytoplankton Manual*. UNESCO, Paris.
- Hernández-Becerril, D.U. and Barón-Campis, S.A. 2008. A new species of the diatom genus *Fryxelliella* (Bacillariophyta), *Fryxelliella pacifica* sp. nov., from the tropical Mexican Pacific. *Phycological Research* 56: 149-155. Doi: <https://doi.org/10.1111/j.1440-1835.2008.00496.x>.
- Hernández-Becerril, D.U., Barón-Campis, S.A. and Ortiz-Lira, H. 2012. A morphological study of epipsammic diatoms (Bacillariophyta) from the tropical Mexican Pacific, including two species of *Aulacodiscus* and a description of the new species *Cerataulus simsae* Hernández-Becerril et Barón-Campis. *Botanica Marina* 55: 73-84. Doi: <https://doi.org/10.1515/bot.2011.124>.
- Hernández-Becerril D.U., Barón-Campis, S.A., Salazar-Paredes, J. and Alonso-Rodríguez, R. 2013. Species of the planktonic diatom genus *Skeletonema* (Bacillariophyta) from the Mexican Pacific Ocean. *Cryptogamie, Algologie* 34: 77-87. Doi: <https://doi.org/10.7872/crya.v34.iss2.2013.77>.
- Hiroimi, J. and Takano, H. 1983. Preliminary note on pelagic marine copepods and their epizoic diatoms. *Bulletin of the Plankton Society of Japan* 30: 171-173.
- Kooistra, W.H.C.F., Gersonde, R., Medlin, L.K. and Mann, D.G. 2007. The origin and evolution of the diatoms: their adaptation to a planktonic existence. In: Falkowsky, P.G. and Knoll A.H. Editors. *Evolution of primary producers in the sea*. Elsevier Academic Press. Cambridge.
- Krasske, G. 1941. Die Kieselalgen des chilenischen Kiistenplanktons. *Archives für Hydrobiologie* 38: 260-287.
- Li, L., Chen C., Sun L., Zhang J., Liang J. and Gao Y. 2020. *Protoraphis* Simonsen, a newly recorded marine epizoic diatom genus for China. *Acta Oceanologica Sinica* 39: 120-126. Doi: <https://doi.org/10.1007/s13131-019-1467-z>.
- Medlin L.K. and Kacmarska I. 2004. Evolution of the diatoms: V. Morphological and cytological support for the major clades and a taxonomic revision. *Phycologia* 43: 245-270. Doi: <https://doi.org/10.2216/i0031-8884-43-3-245.1>.
- Rivera, P.S., González, H.E. and Barrales, H.L. 1986. Cingulum and valve morphology of *Pseudohimantidium* Hustedt & Krasske (Bacillariophyceae). *Phycologia* 25: 19-27. Doi: <https://doi.org/10.2216/i0031-8884-25-1-19.1>
- Round, F.E. 1971. Benthic marine diatoms. *Marine Biology Annual Review* 9: 83-139.
- Round, F.E., Crawford, R.M. and Mann, D.G. 1990. The diatoms. Biology and morphology of the genera. Cambridge University Press, Cambridge.
- Sar, E.A. and Sunesen, I. 2014. The epizoic marine diatom *Sceptronema orientale* (Licmophoraceae, Licmophorales): epitypification and emendation of specific and generic descriptions. *Phytotaxa* 177: 269-279. Doi: <https://doi.org/10.11646/phytotaxa.177.5.3>
- Simonsen, R. 1970. Protoraphidaceae, eine neue Familie der Diatomeen. *Nova Hedwigia, Beihefte* 31: 377-394.
- Skovgaard, A. and Saiz, E. 2006. Seasonal occurrence and role of protistan parasites in coastal marine zooplankton. *Marine Ecology Progress Series* 327: 37-49. Doi: <https://doi.org/10.3354/meps327037>.
- Sunesen, I., Romero, S., Toubes, E. and Sar, E.A. 2015. Morphology and distribution of three araphid diatoms

(Fragilariophyceae, Bacillariophyta) from marine coastal waters of Argentina. *Iheringia, Serie Botanica* 70: 265-78.

Takano, H. 1983. New and rare diatoms from Japanese marine waters-XI. Three new species epizoic on copepods. *Bulletin of the Tokai Regional Fisheries Research Laboratory* 111: 23-35.

Tiffany, M.A. 2011. Epizoic and epiphytic diatoms. In: Seckbach, J. and Kociolek, P., Editors. *The diatom world. Cellular origin, life*

in extreme habitats and astrobiology. Springer, Dordrecht.

Totti, C., Romagnoli, T., de Stefano, M., Di Camillo, C.G. and Bavestrello, G. 2011. The diversity of epizoic diatoms. Relationships between diatoms and marine invertebrates. In: Seckbach, J. and Dubinsky, Z., Editors. *All Flesh Is Grass, Cellular origin, life in extreme habitats and astrobiology* 16. Springer Science+Business Media B.V.

Citar como: Hernández-Becerril, D.U. 2021. Morphology of three marine epizoic diatoms from the Mexican Pacific: *Protoraphis hustediana*, *Pseudohimantidium pacificum* and *Sceptronema orientale*. *Intropica* 16(2): *Postprint*. Doi: <https://doi.org/10.21676/23897864.4066>.